

Arbitrarily Large Finite Quotients Imply No Uniform Bound on Depth

Andrew D. Warshall *
Yale University
Department of Mathematics
10 Hillhouse Avenue
P.O. Box 208283
New Haven, CT 06520-8283
USA

8 February 2006

Abstract

We show that any group with arbitrarily large finite quotients admits generating sets with respect to which it has arbitrarily large finite dead-end depth. This extends a joint result with Riley and partially answers a question asked there.

Let G be any finitely generated group and A a finite generating set for G . Then we define the *depth* (or more verbosely the *dead-end depth*) of an element $g \in G$ with respect to A to be the distance (with respect to the word metric induced by A) from g to the nearest (to g) group element at greater distance than g from the identity. Formally, the depth of g is the distance (in the Cayley graph of G with respect to A) between g and the complement of the radius- $d_A(1, g)$ closed ball about the identity, where d_A refers to distance with respect to A . We define the depth of G with respect to A to be the (possibly infinite) supremum of all the depths of all the elements of G with respect to A .

In general, the depth of a group is not the same for all finite generating sets. In fact, in joint work with Riley [2], we constructed a (finitely *presented*) group which has depth 2 with respect to one generating set and depth ∞ with respect to another. For hyperbolic groups, in contrast, Bogopol'skii [1] has shown that the depth is finite for every generating set. This result suggests the question whether depth is bounded uniformly over all finite generating sets. In [2], we answered this question in the negative for $\mathbb{Z}$, showing that, for all finite n , there exists a generating set for $\mathbb{Z}$ with respect to which it has depth at least n . We here answer it similarly for a broad class of groups, including in particular all indicable groups. Specifically, we prove the following:

*The author acknowledges support under the VIGRE program

Proposition 1. *If G is a finitely generated group with arbitrarily large finite quotients and n is a positive integer then G has a generating set A such that the depth of G with respect to A is $\geq n$.*

This result includes the case where G is indicable, for given $G \twoheadrightarrow \mathbb{Z}$ we have maps $G \twoheadrightarrow C_n$ for arbitrarily large n .

Let G be a finitely generated group with arbitrarily large finite quotients and S a finite generating set for G . We set $a = |S|$.

Proposition 2. *For every integer n' there exists $n \geq n'$, H_n and $\pi_n: G \twoheadrightarrow H_n$ where H_n is a finite group which has an element $h_n \in H_n$ of length n in $(\pi_n(S))^{\pm 1}$ but no element of greater length.*

Proof. Let G_m be a quotient of G , $|G_m| = m$, with α_m being the quotient map $G \twoheadrightarrow G_m$. If G is generated by S then G_m is generated by $\alpha_m(S)$, which consists of a elements, for all m . So there are at most $(2a+1)^n$ words of length $\leq n$ with respect to that generating set, since $(2a+1)^n$ is the number of words of length exactly n in $\alpha_m(S)^{\pm 1} \cup \{1\}$. In particular, there are at most $(2a+1)^n$ group elements within n of the identity with respect to that generating set. Let n be the maximal distance from the identity of any element of G_m ; this must exist since G_m is finite. Then $(2a+1)^n \geq m$, since all group elements are within m of the identity. So if we set $m = (2a+1)^{n'}$ then we have $n \geq n'$. Thus $G_{(2a+1)^{n'}}$ and $\alpha_{(2a+1)^{n'}}$ will serve as H_n and π_n for some $n \geq n'$. $\square$

Notation. We denote $\pi_n(S)$ by T_n .

Proof of Proposition 1. We let N and n' be positive integers and consider $n \geq n'$ and the map $\pi_n: G \twoheadrightarrow H_n$ given by the proposition. Let $A \subset G$ be $B_{S,N}(1) \cap \pi_n^{-1}(T_n)$. I claim that G has depth at least $n \geq n'$ with respect to A . Clearly, A is a generating set so long as $N \geq 1$, since then $S \subseteq A$ since $\pi_n(S) = T_n$. Since h_n is at distance n from the identity with respect to T_n , any $g_n \in \pi_n^{-1}(h_n)$ must be at distance at least n from the identity with respect to $\pi_n^{-1}T_n$, hence *a fortiori* with respect to A .

But I claim that, for every $h \in H_n$, there is an element of $\pi_n^{-1}(h)$ within distance n of the identity with respect to S . To see this, for any $h \in H_n$, take a minimal-length word $\pi_n(s_1)\pi_n(s_2)\dots\pi_n(s_m)$ representing h , $s_i \in S$, $m \leq n$. Then $a = s_1s_2\dots s_m \in \pi_n^{-1}(h)$ is clearly at distance at most $m \leq n$ from the identity. In particular, setting $h = h_n$, there exists $g_n \in \pi_n^{-1}(h_n)$ such that $d_A(1, g_n) = n$. I claim that g_n has depth at least n .

Let $g \in B_{A,d}(g_n)$, where $d < n$. Then g , being the product of g_n , of length n , with up to d words of length N , is, by the triangle inequality, expressible as a word of length at most $n + dN$ in S . Let $k = |\pi_n(g)|$ and divide a geodesic word in S representing G into k roughly equal pieces. Thus $g = u_1u_2\dots u_k$, where the u_i are $\in G$ and their lengths in S are as equal as possible, so that for each u_i $|u_i| < (n + dN)/k + 1$. Let also $t_1t_2\dots t_k$ be a minimal-length word in $T_n^{\pm 1}$ representing $\pi_n(g)$. For $x \in H_n$, we denote by $\phi(x)$ a minimal-length representative of $\pi_n^{-1}(x)$, so by the preceding paragraph we always have

$|\phi(x)| \leq n$. Then we also have

$$\begin{aligned} g = & [u_1 \phi(\pi_n(u_1)^{-1} t_1)] [\phi(\pi_n(u_1)^{-1} t_1)^{-1} u_2 \phi(\pi_n(u_1 u_2)^{-1} t_1 t_2)] \\ & [\phi(\pi_n(u_1 u_2)^{-1} t_1 t_2)^{-1} u_3 \phi(\pi_n(u_1 u_2 u_3)^{-1} t_1 t_2 t_3)] \dots \\ & [\phi(\pi_n(u_1 \dots u_{k-1})^{-1} t_1 \dots t_{k-1})^{-1} u_k]. \end{aligned}$$

We find that, if the v_i are taken to be the factors as indicated by the brackets, g is now expressed as the product $v_1 v_2 \dots v_k$. I claim that all the $v_i \in (\pi_n^{-1}(T_n))^{\pm 1}$. If $i < k$, this is clear since then

$$\pi_n(v_i) = t_{i-1}^{-1} t_{i-2}^{-1} \dots t_1^{-1} \pi_n(u_1 u_2 \dots u_{i-1}) \pi_n(u_i) \pi_n(u_1 u_2 \dots u_i)^{-1} t_1 t_2 \dots t_i = t_i,$$

while if $i = k$ then

$$\begin{aligned} \pi_n(v_i) = \pi_n(v_k) &= t_{k-1}^{-1} t_{k-2}^{-1} \dots t_1^{-1} \pi_n(u_1 u_2 \dots u_{k-1}) \pi_n(u_k) \\ &= t_{k-1}^{-1} t_{k-2}^{-1} \dots t_1^{-1} \pi_n(g) \\ &= t_{k-1}^{-1} t_{k-2}^{-1} \dots t_1^{-1} t_1 t_2 \dots t_k = t_k. \end{aligned}$$

Furthermore, every v_i is obtained from its corresponding u_i by concatenating it with up to two words each of which, being in $\text{im } \phi$, has length at most n , so $|v_i| \leq |u_i| + 2n$. Thus all the v_i will be $\in A^{\pm 1}$ so long as all the $|u_i| + 2n \leq N$, which will occur if $(n + dN)/k + 1 + 2n \leq N$, for which (since $k \geq n - d$ by the triangle inequality) it suffices that $(n + dN)/(n - d) + 2n + 1 \leq N$, which will be true so long as $n > 2d$ and $N \geq (2n^2 + 2nd + 2n - d)/(n - 2d)$. $\square$

References

- [1] O.V. Bogopol'skii, Infinite commensurable hyperbolic groups are bi-Lip-schitz equivalent, *Algebra and Logic*, 36(3):155–63, 1997.
- [2] T.R. Riley and A.D. Warshall, The unbounded dead-end depth property is not a group invariant, to appear in *International Journal of Algebra and Computation*, [arXiv:math.GR/0504121](https://arxiv.org/abs/math.GR/0504121).